

Drug Policy

Policy:	230201	Initial Effective Date: 3/31/2023
Code(s):	HCPCS J3590	Annual Review Date: 02/20/2024
SUBJECT:	Briumvi™ (ublituximab-xiij)	Last Revised Date: 02/20/2024

Subject to Site of Care

Prior approval is required for some or all procedure codes listed in this Corporate Drug Policy.

Initial and renewal requests for the medication(s) listed in this policy are subject to site of care management. When billed under the medical benefit, administration of the medication will be restricted to a non-hospital facility-based location (i.e., home infusion provider, provider’s office, free-standing ambulatory infusion center) unless the member meets the site of care exception criteria. To view the exception criteria and a list of medications subject to site of care management please [click here](#).

POLICY STATEMENT

This policy involves the use of Briumvi. Prior authorization is recommended for pharmacy and medical benefit coverage of Briumvi. Approval is recommended for those who meet the conditions of coverage in the **Criteria, Dosing, and Initial/Extended Approval** for the diagnosis provided. Requests for uses not listed in this policy will be reviewed for evidence of efficacy and for medical necessity on a case-by-case basis.

Because of the specialized skills required for evaluation and diagnosis of patients treated with Briumvi as well as the monitoring required for AEs and long-term efficacy, initial approval requires Briumvi be prescribed by or in consultation with a physician who specializes in the condition being treated. All approvals for initial therapy are provided for the initial approval duration noted below; if reauthorization is allowed, a response to therapy is required for continuation of therapy unless otherwise noted below.

RECOMMENDED AUTHORIZATION CRITERIA

Coverage of Briumvi is recommended in those who meet the following criteria:

FDA-Approved Indication

1. **Multiple Sclerosis, Relapsing Forms.** Approve 1 year if the patient meets one of the following (A or B):
 - A) **Initial Therapy.** Approve if the patient meets all the following (i, ii, iii, and iv):
 - i. Patient is \geq 18 years of age; AND
 - ii. Patient has a relapsing form of multiple sclerosis; AND

This document is subject to the disclaimer found at https://provider.medmutual.com/tools_and_resources/Care_Management/MedPolicies/Disclaimer.aspx and is subject to change. Always verify with the most current version at https://provider.medmutual.com/tools_and_resources/Care_Management/MedPolicies/Disclaimer.aspx or https://provider.medmutual.com/TOOLS_and_RESOURCES/Care_Management/ExpressScripts.aspx.

Drug Policy

Note: Examples of relapsing forms of multiple sclerosis include clinically isolated syndrome, relapsing remitting disease, and active secondary progressive disease.

- iii. According to the prescriber, the patient has experienced inadequate efficacy or significant intolerance to one disease-modifying agent used for multiple sclerosis; AND

Note: See [Appendix](#) for examples.

- iv. Medication is prescribed by or in consultation with a neurologist or a physician who specializes in the treatment of multiple sclerosis; OR

- B) Patient is Currently Receiving Briumvi for \geq 1 Year.** Approve if the patient meets all of the following criteria (i, ii, iii and iv):

Note: A patient who has received < 1 year of therapy or who is restarting therapy with Briumvi should be considered under criterion 1A (Multiple Sclerosis [Relapsing Forms], Initial Therapy).

- i. Patient is \geq 18 years of age; AND
ii. Patient has a relapsing form of multiple sclerosis; AND

Note: Examples of relapsing forms of multiple sclerosis include clinically isolated syndrome, relapsing remitting disease, and active secondary progressive disease.

- iii. Patient meets one of the following [(1) or (2)]:

- (1) Patient experienced a beneficial clinical response when assessed by at least one objective measure; OR

Note: Examples include stabilization or reduced worsening in disease activity as evaluated by magnetic resonance imaging (MRI) [absence or a decrease in gadolinium enhancing lesions, decrease in the number of new or enlarging T2 lesions]; stabilization or reduced worsening on the Expanded Disability Status Scale (EDSS) score; achievement in criteria for No Evidence of Disease Activity-3 (NEDA-3) or NEDA-4; improvement on the fatigue symptom and impact questionnaire-relapsing multiple sclerosis (FSIQ-RMS) scale; reduction or absence of relapses; improvement or maintenance on the six-minute walk test or 12-Item Multiple Sclerosis Walking Scale; improvement on the Multiple Sclerosis Functional Composite (MSFC) score; and/or attenuation of brain volume loss; OR

- (2) Patient experienced stabilization, slowed progression, or improvement in at least one symptom such as motor function, fatigue, vision, bowel/bladder function, spasticity, walking/gait, or pain/numbness/tingling sensation; AND

- iv. Medication is prescribed by or in consultation with a neurologist or a physician who specializes in the treatment of multiple sclerosis.

Dosing in Multiple Sclerosis, Relapsing Forms. Dosing must meet the following (medical benefit only):

Initial dosing:

- First Infusion: 150 mg intravenous infusion
- Second Infusion: 450 mg intravenous infusion administered two weeks after the first infusion

Subsequent doses:

450 mg intravenous infusion administered 24 weeks after the **first** infusion and every 24 weeks thereafter

Initial Approval/ Extended Approval.

This document is subject to the disclaimer found at https://provider.medmutual.com/tools_and_resources/Care_Management/MedPolicies/Disclaimer.aspx and is subject to change. Always verify with the most current version at https://provider.medmutual.com/tools_and_resources/Care_Management/MedPolicies/Disclaimer.aspx or https://provider.medmutual.com/TOOLS_and_RESOURCES/Care_Management/ExpressScripts.aspx.

Drug Policy

- A) *Initial Approval*: 1 year
- B) *Extended Approval*: 1 year

CONDITIONS NOT RECOMMENDED FOR APPROVAL

Briumvi has not been shown to be effective, or there are limited or preliminary data or potential safety concerns that are not supportive of general approval for the following conditions. (Note: This is not an exhaustive list of Conditions Not Recommended for Approval).

1. **Concurrent Use with Other Disease-Modifying Agents Used for Multiple Sclerosis.** These agents are not indicated for use in combination (See [Appendix](#) for examples). Additional data are required to determine if use of disease-modifying multiple sclerosis agents in combination is safe and provides added efficacy.
2. Coverage is not recommended for circumstances not listed in the Recommended Authorization Criteria. Criteria will be updated as new published data are available.

Documentation Requirements:

The Company reserves the right to request additional documentation as part of its coverage determination process. The Company may deny reimbursement when it has determined that the drug provided or services performed were not medically necessary, investigational or experimental, not within the scope of benefits afforded to the member and/or a pattern of billing or other practice has been found to be either inappropriate or excessive. Additional documentation supporting medical necessity for the services provided must be made available upon request to the Company. Documentation requested may include patient records, test results and/or credentials of the provider ordering or performing a service. The Company also reserves the right to modify, revise, change, apply and interpret this policy at its sole discretion, and the exercise of this discretion shall be final and binding.

REFERENCES

1. Briumvi® intravenous infusion [prescribing information]. Morrisville, NC: TG Therapeutics; December 2022.
2. A Consensus Paper by the Multiple Sclerosis Coalition. The use of disease-modifying therapies in multiple sclerosis. Updated September 2019. Available at: https://www.nationalmssociety.org/NationalMSSociety/media/MSNationalFiles/Brochures/DMT_Consensus_MS_Coalition.pdf. Accessed on November 14, 2023.
3. McGinley MP, Goldschmidt C, Rae-Grant AD. Diagnosis and treatment of multiple sclerosis. A review. *JAMA*. 2021;325(8):765-779.
4. The Medical Letter on Drugs and Therapeutics. Drugs for multiple sclerosis. *Med Lett Drugs Ther*. 2021;63(1620):42-48.
5. Lublin FD, Reingold SC, Cohen JA, et al. Defining the clinical course of multiple sclerosis: the 2013 revisions. *Neurology*. 2014;83:278-286.
6. Thompson AJ, Banwell BL, Barkhof F, et al. Diagnosis of multiple sclerosis: 2017 revisions of the McDonald criteria. *Lancet Neurol*. 2018;17(2):162-173.

This document is subject to the disclaimer found at https://provider.medmutual.com/tools_and_resources/Care_Management/MedPolicies/Disclaimer.aspx and is subject to change. Always verify with the most current version at https://provider.medmutual.com/tools_and_resources/Care_Management/MedPolicies/Disclaimer.aspx or https://provider.medmutual.com/TOOLS_and_RESOURCES/Care_Management/ExpressScripts.aspx.

Drug Policy

APPENDIX

Medication	Mode of Administration
Aubagio® (teriflunomide tablets)	Oral
Avonex® (interferon beta-1a intramuscular injection)	Injection (self-administered)
Bafiertam® (monomethyl fumarate delayed-release capsules)	Oral
Betaseron® (interferon beta-1b subcutaneous injection)	Injection (self-administered)
Briumvi™ (ublituximab-xijj intravenous infusion)	Intravenous infusion
Copaxone® (glatiramer acetate subcutaneous injection, generic)	Injection (self-administered)
Extavia® (interferon beta-1b subcutaneous injection)	Injection (self-administered)
Gilenya® (fingolimod capsules, generic)	Oral
Glatopa® (glatiramer acetate subcutaneous injection)	Injection (self-administered)
Kesimpta® (ofatumumab subcutaneous injection)	Injection (self-administered)
Lemtrada® (alemtuzumab intravenous infusion)	Intravenous infusion
Mavenclad® (cladribine tablets)	Oral
Mayzent® (siponimod tablets)	Oral
Ocrevus® (ocrelizumab intravenous infusion)	Intravenous infusion
Plegridy® (peginterferon beta-1a subcutaneous or intramuscular injection)	Injection (self-administered)
Ponvory™ (ponesimod tablets)	Oral
Rebif® (interferon beta-1a subcutaneous injection)	Injection (self-administered)
Tascenso ODT™ (fingolimod orally disintegrating tablets)	Oral
Tecfidera® (dimethyl fumarate delayed-release capsules, generic)	Oral
Tysabri® (natalizumab intravenous infusion)	Intravenous infusion
Vumerity® (diroximel fumarate delayed-release capsules)	Oral
Zeposia® (ozanimod capsules)	Oral

FOR MEDICAL BENEFIT COVERAGE REQUESTS:

Prior approval is required for HCPCS Codes J3590

†When *unclassified biologics* (J3590) is determined to be Briumvi

Edits and Denials:

Prior approval: Prior approval is required for Briumvi (**HCPCS Codes J3590**). Requests for prior approval will be authorized by a nurse reviewer if submitted documentation meets criteria outlined within the Corporate Medical Policy.

This document is subject to the disclaimer found at https://provider.medmutual.com/tools_and_resources/Care_Management/MedPolicies/Disclaimer.aspx and is subject to change. Always verify with the most current version at https://provider.medmutual.com/tools_and_resources/Care_Management/MedPolicies/Disclaimer.aspx or https://provider.medmutual.com/TOOLS_and_RESOURCES/Care_Management/ExpressScripts.aspx.

Drug Policy

Requests for prior approval will be forwarded to a qualified physician reviewer if submitted documentation does not meet criteria outlined within Corporate Medical Policy.

TOPPS: Claims received with **HCPCS Codes J3590** will pend with **Remark Code M3M or M4M** and will be adjudicated in accordance with the Corporate Medical Policy.

Liability: A participating provider will be required to write off charges denied as not medically necessary.

HCPCS Code(s):	
J3590	Unclassified biologics

This document is subject to the disclaimer found at https://provider.medmutual.com/tools_and_resources/Care_Management/MedPolicies/Disclaimer.aspx and is subject to change. Always verify with the most current version at https://provider.medmutual.com/tools_and_resources/Care_Management/MedPolicies/Disclaimer.aspx or https://provider.medmutual.com/TOOLS_and_RESOURCES/Care_Management/ExpressScripts.aspx.