

# Drug Policy

<b>Policy:</b>	<b>Bimzelx® (bimekizuma-bkzx subcutaneous injection)</b>	<b>Annual Review Date:</b> <b>8/21/2025</b> <b>Last Revised Date:</b> <b>8/21/2025</b>
----------------	--	---

## OVERVIEW

Bimzelx, an interleukin (IL)-17A and IL-17F blocker, is indicated for treatment of adults with moderate to severe plaque psoriasis, ankylosing spondylitis, psoriatic arthritis and non-radiographic spondyloarthritis.

## POLICY STATEMENT

policy involves the use of Bimzelx. Prior authorization is recommended for pharmacy benefit coverage of Bimzelx. Approval is recommended for those who meet the conditions of coverage in the **Criteria and Initial/Extended Approval** for the diagnosis provided. **Conditions Not Recommended for Approval** are listed following the recommended authorization criteria. Requests for uses not listed in this policy will be reviewed for evidence of efficacy and for medical necessity on a case-by-case basis.

Because of the specialized skills required for evaluation and diagnosis of patients treated with Bimzelx as well as the monitoring required for adverse events and long-term efficacy, initial approval requires Bimzelx be prescribed by or in consultation with a physician who specializes in the condition being treated. All approvals for initial therapy are provided for the initial approval duration noted below; if reauthorization is allowed, a response to therapy is required for continuation of therapy unless otherwise noted below.

## RECOMMENDED AUTHORIZATION CRITERIA

Coverage of Bimzelx is recommended in those who meet the following criteria:

### FDA-Approved Indication

1. **Ankylosing Spondylitis.** Approve for the duration noted if the patient meets ONE of the following (A or B):
  - A) **Initial Therapy.** Approve for 6 months if the patient meets ALL of the following (i, ii, iii, iv, and v):
    - i. Patient is  $\geq$  18 years of age; AND
    - ii. The prescriber attests that the patient has been assessed and evaluated for risks of suicidal ideation or behavior versus benefits of therapy; AND
    - iii. The patient does **not** have moderately severe to severe depression; AND
    - iv. Within the past 5 years, the patient does **not** have a history of suicidal ideation or suicidal behavior; AND
    - v. The medication is prescribed by or in consultation with a rheumatologist.
  - B) **Patient is Currently Receiving Bimzelx.** Approve for 1 year if the patient meets ALL of the following (i, ii, iii, iv, and v):

This document is subject to the disclaimer found at <https://www.medmutual.com/For-Providers/Policies-and-Standards/CorporateMedicalDisclaimer.aspx> and is subject to change. <https://www.medmutual.com/For-Providers/Policies-and-Standards/Prescription-Drug-Resources.aspx>

# Drug Policy

- i. Patient has been established on therapy for at least 6 months; AND  
Note: A patient who has received < 6 months of therapy or who is restarting therapy is reviewed under criterion A (Initial Therapy).
- ii. The prescriber attests that the patient has been assessed and evaluated for risks of suicidal ideation or behavior versus benefits of therapy; AND
- iii. The patient does **not** have moderately severe to severe depression; AND
- iv. According to the prescriber, the patient does **not** have suicidal ideation or suicidal behavior; AND
- v. Patient meets at least ONE of the following (a or b):
  - a) When assessed by at least one objective measure, patient experienced a beneficial clinical response from baseline (prior to initiating Bimzelx); OR  
Note: Examples of objective measures include Ankylosing Spondylitis Disease Activity Score (ASDAS), Ankylosing Spondylitis Quality of Life Scale (ASQoL), Bath Ankylosing Spondylitis Disease Activity Index (BASDAI), Bath Ankylosing Spondylitis Functional Index (BASFI), Bath Ankylosing Spondylitis Global Score (BAS-G), Bath Ankylosing Spondylitis Metrology Index (BASMI), Dougados Functional Index (DFI), Health Assessment Questionnaire for the Spondylarthropathies (HAQ-S), and/or serum markers (e.g., C-reactive protein, erythrocyte sedimentation rate).
  - b) Compared with baseline (prior to initiating Bimzelx), patient experienced an improvement in at least one symptom, such as decreased pain or stiffness, or improvement in function or activities of daily living.

## 2. **Hidradenitis Suppurativa.** Approve for the duration noted if the patient meets ONE of the following (A or B):

- A) Initial Therapy. Approve for 3 months if the patient meets ALL of the following (i, ii, iii, iv, v, and vi):
  - i. Patient is ≥ 18 years of age; AND
  - ii. Patient has tried at least one other therapy; AND  
Note: Examples include intralesional or oral corticosteroids (e.g., triamcinolone, prednisone), systemic antibiotics (e.g., clindamycin, dicloxacillin, erythromycin), and isotretinoin.
  - iii. According to the prescriber, the patient has been evaluated for risks of suicidal ideation or behavior versus benefits of therapy; AND
  - iv. According to the prescriber, the patient does **not** have moderately severe to severe depression; AND
  - v. According to the prescriber, within the past 5 years, the patient does **not** have a history of suicidal ideation or suicidal behavior; AND
  - vi. The medication is prescribed by or in consultation with a dermatologist; OR
- B) Patient is Currently Receiving Bimzelx. Approve for 1 year if the patient meets ALL of the following (i, ii, iii, iv, v, and vi):
  - i. Patient has been established on Bimzelx for at least 3 months; AND  
Note: A patient who has received < 3 months of therapy or who is restarting therapy with Bimzelx is reviewed under criterion A (Initial Therapy).
  - ii. According to the prescriber, the patient has been evaluated for risks of suicidal ideation or behavior versus benefits of therapy; AND
  - iii. According to the prescriber, the patient does **not** have moderately severe to severe depression; AND
  - iv. According to the prescriber, the patient does **not** have suicidal ideation or suicidal behavior; AND
  - v. When assessed by at least one objective measure, patient experienced a beneficial clinical response from baseline (prior to initiating Bimzelx); AND

# Drug Policy

**Note:** Examples of objective measures include Hurley staging, Sartorius score, Physician Global Assessment, and Hidradenitis Suppurativa Severity Index.

- vi. Compared with baseline (prior to initiating Bimzelx), patient experienced an improvement in at least one symptom, such as decreased pain or drainage of lesions, nodules, or cysts.

**3. Non-Radiographic Axial Spondyloarthritis.** Approve for the duration noted if the patient meets ONE of the following (A or B):

**A) Initial Therapy.** Approve for 6 months if the patient meets ALL of the following (i, ii, iii, iv, v, and vi):

- i. Patient is  $\geq 18$  years of age; AND
- ii. Patient has objective signs of inflammation, defined as at least ONE of the following (a or b):
  - a) C-reactive protein elevated beyond the upper limit of normal for the reporting laboratory; OR
  - b) Sacroiliitis reported on magnetic resonance imaging; AND
- iii. The prescriber attests that the patient has been assessed and evaluated for risks of suicidal ideation or behavior versus benefits of therapy; AND
- iv. The patient does **not** have moderately severe to severe depression; AND
- v. Within the past 5 years, the patient does **not** have a history of suicidal ideation or suicidal behavior; AND
- vi. The medication is prescribed by or in consultation with a rheumatologist.

**B) Patient is Currently Receiving Bimzelx.** Approve for 1 year if the patient meets ALL of the following (i, ii, iii, iv, and v):

- i. Patient has been established on therapy for at least 6 months; AND  
**Note:** A patient who has received < 6 months of therapy or who is restarting therapy is reviewed under criterion A (Initial Therapy).
- ii. The prescriber attests that the patient has been assessed and evaluated for risks of suicidal ideation or behavior versus benefits of therapy; AND
- iii. The patient does **not** have moderately severe to severe depression; AND
- iv. According to the prescriber, the patient does **not** have suicidal ideation or suicidal behavior; AND
- v. Patient meets at least ONE of the following (a or b):
  - a) When assessed by at least one objective measure, patient experienced a beneficial clinical response from baseline (prior to initiating Bimzelx); OR  
**Note:** Examples of objective measures include Ankylosing Spondylitis Disease Activity Score (ASDAS), Ankylosing Spondylitis Quality of Life Scale (ASQoL), Bath Ankylosing Spondylitis Disease Activity Index (BASDAI), Bath Ankylosing Spondylitis Functional Index (BASFI), Bath Ankylosing Spondylitis Global Score (BAS-G), Bath Ankylosing Spondylitis Metrology Index (BASMI), Dougados Functional Index (DFI), Health Assessment Questionnaire for the Spondylarthropathies (HAQ-S), and/or serum markers (e.g., C-reactive protein, erythrocyte sedimentation rate).
  - b) Compared with baseline (prior to initiating Bimzelx), patient experienced an improvement in at least one symptom, such as decreased pain or stiffness, or improvement in function or activities of daily living.

**4. Plaque Psoriasis.** Approve for the duration noted if the patient meets ONE of the following (A or B):

**C) Initial Therapy.** Approve for 3 months if the patient meets ALL of the following (i, ii, iii, iv, v, and vi):

- i. Patient is  $\geq 18$  years of age; AND
- ii. Patient meets ONE of the following (a or b):

# Drug Policy

- a) Patient has tried at least one traditional systemic agent for psoriasis for at least 3 months, unless intolerant; OR

Note: Examples include methotrexate, cyclosporine, or acitretin. A 3-month trial of psoralen plus ultraviolet A light (PUVA) also counts. An exception to the requirement for a trial of one traditional systemic agent for psoriasis can be made if the patient has already had a 3-month trial or previous intolerance to at least one biologic other than the requested drug. A biosimilar of the requested biologic does not count. Refer to [Appendix](#) for examples of biologics used for plaque psoriasis. A patient who has already tried a biologic for psoriasis is not required to “step back” and try a traditional systemic agent for psoriasis.

- b) Patient has a contraindication to methotrexate, as determined by the prescriber; AND

- iii. The prescriber attests that the patient has been assessed and evaluated for risks of suicidal ideation or behavior versus benefits of therapy; AND

- iv. The patient does not have moderately severe to severe depression; AND

- v. Within the past 5 years, the patient does not have a history of suicidal ideation or suicidal behavior; AND

- vi. The medication is prescribed by or in consultation with a dermatologist.

- D) Patient is Currently Receiving Bimzelx. Approve for 1 year if the patient meets ALL of the following (i, ii, iii, iv, v, and vi):

- i. Patient has been established on therapy for at least 3 months; AND

Note: A patient who has received < 3 months of therapy or who is restarting therapy is reviewed under criterion A (Initial Therapy).

- ii. The prescriber attests that the patient has been assessed and evaluated for risks of suicidal ideation or behavior versus benefits of therapy; AND

- iii. The patient does not have moderately severe to severe depression; AND

- iv. According to the prescriber, the patient does not have suicidal ideation or suicidal behavior; AND

- v. Patient experienced a beneficial clinical response, defined as improvement from baseline (prior to initiating Bimzelx) in at least one of the following: estimated body surface area, erythema, induration/thickness, and/or scale of areas affected by psoriasis; AND

- vi. Compared with baseline (prior to receiving Bimzelx), patient experienced an improvement in at least one symptom, such as decreased pain, itching, and/or burning.

- 5. **Psoriatic Arthritis.** Approve for the duration noted if the patient meets ONE of the following (A or B):

- A) Initial Therapy. Approve for 6 months if the patient meets ALL of the following (i, ii, iii, iv, and v):

- i. Patient is  $\geq 18$  years of age; AND

- ii. The prescriber attests that the patient has been assessed and evaluated for risks of suicidal ideation or behavior versus benefits of therapy; AND

- iii. The patient does not have moderately severe to severe depression; AND

- iv. Within the past 5 years, the patient does not have a history of suicidal ideation or suicidal behavior; AND

- v. The medication is prescribed by or in consultation with a rheumatologist or a dermatologist.

- B) Patient is Currently Receiving Bimzelx. Approve for 1 year if the patient meets ALL of the following (i, ii, iii, iv, and v):

- i. Patient has been established on therapy for at least 6 months; AND

# Drug Policy

Note: A patient who has received < 6 months of therapy or who is restarting therapy is reviewed under criterion A (Initial Therapy).

- ii. The prescriber attests that the patient has been assessed and evaluated for risks of suicidal ideation or behavior versus benefits of therapy; AND
- iii. The patient does **not** have moderately severe to severe depression; AND
- iv. According to the prescriber, the patient does **not** have suicidal ideation or suicidal behavior; AND
- v. Patient meets at least ONE of the following (a or b):
  - a) When assessed by at least one objective measure, patient experienced a beneficial clinical response from baseline (prior to initiating Bimzelx); OR  
Note: Examples of standardized measures of disease activity include Disease Activity Index for Psoriatic Arthritis (DAPSA), Composite Psoriatic Disease Activity Index (CPDAI), Psoriatic Arthritis Disease Activity Score (PsA DAS), Grace Index, Leeds Enthesitis Score (LEI), Spondyloarthritis Consortium of Canada (SPARCC) enthesitis score, Leeds Dactylitis Instrument Score, Minimal Disease Activity (MDA), Psoriatic Arthritis Impact of Disease (PsAID-12), and/or serum markers (e.g., C-reactive protein, erythrocyte sedimentation rate).
  - b) Compared with baseline (prior to initiating Bimzelx), patient experienced an improvement in at least one symptom, such as less joint pain, morning stiffness, or fatigue; improved function or activities of daily living; decreased soft tissue swelling in joints or tendon sheaths.

---

## CONDITIONS NOT RECOMMENDED FOR APPROVAL

Bimzelx has not been shown to be effective, or there are limited or preliminary data or potential safety concerns that are not supportive of general approval for the following conditions. (Note: This is not an exhaustive list of Conditions Not Recommended for Approval).

1. **Concurrent Use with other Biologics or with Targeted Synthetic Disease-Modifying Antirheumatic Drugs (DMARDs).** Bimzelx should not be administered in combination with a biologic used for an inflammatory condition (see [Appendix](#) for examples). Combination therapy with biologics and/or biologics + targeted synthetic DMARDs has a potential for a higher rate of adverse effects and lacks controlled trial data in support of additive efficacy.  
Note: This does NOT exclude the use of conventional synthetic DMARDs (e.g., methotrexate, leflunomide, hydroxychloroquine, and sulfasalazine) in combination with Bimzelx.
2. **Inflammatory Bowel Disease (i.e., Crohn's disease, ulcerative colitis).** Exacerbations of inflammatory bowel disease, in some cases serious, occurred in clinical trials involving patients treated with Bimzelx.<sup>1</sup>
3. Coverage is not recommended for circumstances not listed in the Recommended Authorization Criteria. Criteria will be updated as new published data are available.

## Documentation Requirements:

---

This document is subject to the disclaimer found at <https://www.medmutual.com/For-Providers/Policies-and-Standards/CorporateMedicalDisclaimer.aspx> and is subject to change. <https://www.medmutual.com/For-Providers/Policies-and-Standards/Prescription-Drug-Resources.aspx>



# Drug Policy

The Company reserves the right to request additional documentation as part of its coverage determination process. The Company may deny reimbursement when it has determined that the drug provided or services performed were not medically necessary, investigational, or experimental, not within the scope of benefits afforded to the member and/or a pattern of billing or other practice has been found to be either inappropriate or excessive. Additional documentation supporting medical necessity for the services provided must be made available upon request to the Company. Documentation requested may include patient records, test results and/or credentials of the provider ordering or performing a service. The Company also reserves the right to modify, revise, change, apply and interpret this policy at its sole discretion, and the exercise of this discretion shall be final and binding.

## REFERENCES

1. Bimzelx® subcutaneous injection [prescribing information]. Smyrna, GA: UCB; October 2023.
2. Menter A, Strober BE, Kaplan DH, et al. Joint AAD-NPF guidelines of care for the management and treatment of psoriasis with biologics. *J Am Acad Dermatol*. 2019 80(4):1029-1072.

## APPENDIX

	Mechanism of Action	Examples of Inflammatory Indications*
<b>Biologics</b>		
<b>Adalimumab SC Products</b> (Humira®, biosimilars)	Inhibition of TNF	AS, CD, JIA, PsO, PsA, RA, UC
<b>Cimzia®</b> (certolizumab pegol SC injection)	Inhibition of TNF	AS, CD, nr-axSpA, PsO, PsA, RA
<b>Etanercept SC Products</b> (Enbrel®, biosimilars)	Inhibition of TNF	AS, JIA, PsO, PsA
<b>Zymfentra®</b> (infliximab-dyyb SC injection)	Inhibition of TNF	CD, UC
<b>Infliximab IV Products</b> (Remicade®, biosimilars)	Inhibition of TNF	AS, CD, PsO, PsA, RA, UC
<b>Simponi®, Simponi® Aria™</b> (golimumab SC injection, golimumab IV infusion)	Inhibition of TNF	SC formulation: AS, PsA, RA, UC
		IV formulation: AS, PJIA, PsA, RA
<b>Actemra®</b> (tocilizumab IV infusion, tocilizumab SC injection)	Inhibition of IL-6	SC formulation: PJIA, RA, SJIA
		IV formulation: PJIA, RA, SJIA
<b>Kevzara®</b> (sarilumab SC injection)	Inhibition of IL-6	RA, PMR
<b>Orencia®</b> (abatacept IV infusion, abatacept SC injection)	T-cell costimulation modulator	SC formulation: JIA, PSA, RA
		IV formulation: JIA, PsA, RA
<b>Rituximab IV Products</b> (Rituxan®, biosimilars)	CD20-directed cytolytic antibody	RA
<b>Kineret®</b> (anakinra SC injection)	Inhibition of IL-1	JIA*, RA
<b>Stelara®</b> (ustekinumab SC injection, ustekinumab IV infusion)	Inhibition of IL-12/23	SC formulation: CD, PsO, PsA, UC
		IV formulation: CD, UC
<b>Siliq™</b> (brodalumab SC injection)	Inhibition of IL-17RA	PsO
<b>Bimzelx®</b> (bimekizumab-bkzx SC injection)	Inhibition of IL-17A and IL-17F	PsO
<b>Cosentyx®</b> (secukinumab SC injection, secukinumab IV infusion)	Inhibition of IL-17A	SC formulation: AS, ERA, nr-axSpA, PsO, PsA
		IV formulation: AS, nr-axSpA, PsA
<b>Taltz®</b> (ixekizumab SC injection)	Inhibition of IL-17A	AS, nr-axSpA, PsO, PsA
<b>Ilumya™</b> (tildrakizumab-asnm SC injection)	Inhibition of IL-23	PsO
<b>Skyrizi®</b> (risankizumab-rzaa SC injection, risankizumab-rzaa IV infusion)	Inhibition of IL-23	SC formulation: CD, PSA, PsO
		IV formulation: CD
<b>Tremfya™</b> (guselkumab SC injection)	Inhibition of IL-23	PsO

This document is subject to the disclaimer found at <https://www.medmutual.com/For-Providers/Policies-and-Standards/CorporateMedicalDisclaimer.aspx> and is subject to change. <https://www.medmutual.com/For-Providers/Policies-and-Standards/Prescription-Drug-Resources.aspx>

# Drug Policy

<b>Entyvio™</b> (vedolizumab IV infusion, vedolizumab SC injection)	Integrin receptor antagonist	SC formulation: UC IV formulation: CD, UC
<b>Oral Therapies/Targeted Synthetic DMARDs</b>		
<b>Otezla®</b> (apremilast tablets)	Inhibition of PDE4	PsO, PsA
<b>Cibinqo™</b> (abrocitinib tablets)	Inhibition of JAK pathways	AD
<b>Olumiant®</b> (baricitinib tablets)	Inhibition of JAK pathways	RA
<b>Rinvoq®</b> (upadacitinib extended-release tablets)	Inhibition of JAK pathways	AD, AS, nr-axSpA, RA, PsA, UC
<b>Sotyktu™</b> (deucravacitinib tablets)	Inhibition of TYK2	PsO
<b>Xeljanz®</b> (tofacitinib tablets)	Inhibition of JAK pathways	RA, PJIA, PsA, UC
<b>Xeljanz® XR</b> (tofacitinib extended-release tablets)	Inhibition of JAK pathways	RA, PsA, UC

\* Not an all-inclusive list of indications (e.g., oncology indications and rare inflammatory conditions are not listed). Refer to the prescribing information for the respective agent for FDA-approved indications; SC – Subcutaneous; TNF – Tumor necrosis factor; AS – Ankylosing spondylitis; CD – Crohn’s disease; JIA – Juvenile idiopathic arthritis; PsO – Plaque psoriasis; PsA – Psoriatic arthritis; RA – Rheumatoid arthritis; UC – Ulcerative colitis; nr-axSpA – Non-radiographic axial spondyloarthritis; IV – Intravenous, PJIA – Polyarticular juvenile idiopathic arthritis; IL – Interleukin; SJIA – Systemic juvenile idiopathic arthritis; PMR – Polymyalgia rheumatic; ^ Off-label use of Kineret in JIA supported in guidelines; ERA – Enthesitis-related arthritis; DMARD – Disease-modifying antirheumatic drug; PDE4 – Phosphodiesterase 4; JAK – Janus kinase; AD – Atopic dermatitis; TYK2 – Tyrosine kinase 2.