

Drug Policy

Policy: Impacted Drugs:	Proton Pump Inhibitors (PPI) Preferred Step Therapy Program <ul style="list-style-type: none"> • Dexlansoprazole • Esomeprazole oral packets • Lansoprazole orally disintegrating tablets • Pantoprazole oral packets • Voquezna 	Annual Review Date: 01/16/2025 Last Revised Date: 01/16/2025
--	--	---

OVERVIEW

Proton pump inhibitors (PPIs) [i.e., esomeprazole, dexlansoprazole, lansoprazole, omeprazole, pantoprazole, and rabeprazole] are commonly used antisecretory agents that are highly effective at suppressing gastric acid and subsequently treating associated conditions, including gastroesophageal reflux disease (GERD).

POLICY STATEMENT

A step therapy program has been developed to encourage the use of one generic Step 1 product prior to the use of a Step 2 product. If the step therapy rule is not met for a Step 2 agent at the point of service, coverage will be determined by the step therapy criteria below. All approvals are provided for 1 year in duration. **Please note** that not all plans cover over-the-counter drugs, please check corresponding benefit materials for additional information.

Preferred Medications (step 1)

- Omeprazole delayed release capsules and tablets (generic, Rx and OTC)
- Pantoprazole delayed release tablets (generic)

Non-Preferred Medications (step 2)

- Dexlansoprazole delayed release capsule (generic)
- Esomeprazole oral packets (generic)
- Lansoprazole orally disintegrating tablets (generic, Rx and OTC)
- Pantoprazole oral packets (generic)
- Voquezna tablets

Automation: Patients with a history of one Step 1 drug within the 130-day look-back period are excluded from step therapy.

PREFERRED STEP THERAPY CRITERIA

This document is subject to the disclaimer found at <https://www.medmutual.com/For-Providers/Policies-and-Standards/CorporateMedicalDisclaimer.aspx> and is subject to change. <https://www.medmutual.com/For-Providers/Policies-and-Standards/Prescription-Drug-Resources.aspx>

Drug Policy

1. If the patient has tried a Step 1 PPI under the supervision of a physician, then authorization may be given for a Step 2 PPI product. **Note:** A trial of a generic OTC PPI would qualify if OTC PPIs are a covered benefit and the patient was using it for at least 14 days.
2. If the patient is < 1 year of age OR has difficulty swallowing, then authorization may be given for generic esomeprazole delayed release granules for oral suspension (packets), lansoprazole orally disintegrating tablets, or pantoprazole delayed release oral suspension (packets). **Note:** Esomeprazole is indicated down to 1 month of age.

Initial Approval/ Extended Approval.

A) *Initial Approval:* 2 years

B) *Extended Approval:* 2 years

Step Therapy Exception Criteria

In certain situations, the patient is not required to trial preferred agents. Approve for 1 year if the patient meets the following (A, B, or C):

- A. The patient has an atypical diagnosis and/or unique patient characteristics which prevent use of all preferred agents. If so, please list diagnosis and/or patient characteristics*; **OR**
- B. The patient has a contraindication to all preferred agents. If so, please list the contraindications to each preferred agent*; **OR**
- C. The patient is continuing therapy with the requested non-preferred agent after being stable for at least 90 days [verification in prescription claims history required] or, if not available, [verification by prescribing physician required] AND meets ONE of the following:
 1. The patient has at least 130 days of prescription claims history on file and claims history supports that the patient has received the requested non-preferred agent for 90 days within a 130-day look-back period AND there is no generic equivalent available for the requested nonpreferred product (i.e. AA-rated or AB-rated to the requested nonpreferred product); **OR**
 2. When 130 days of the patient's prescription claims history file is unavailable for verification, the prescriber must verify that the patient has been receiving the requested non-preferred agent for 90 days AND that the patient has been receiving the requested non-preferred agent via paid claims (i.e. the patient has NOT been receiving samples or coupons or other types of waivers in order to obtain access to the requested non-preferred agent) AND there is no generic equivalent available for the requested nonpreferred product (i.e. AA-rated or AB-rated to the requested nonpreferred product).

***Documentation Required:** When documentation is required, the prescriber must provide written documentation supporting the trials of these other agents, noted in the criteria as (*). Documentation should include chart notes, prescription claims records, and/or prescription receipts.

Approval Duration: All approvals for continuation of therapy are provided for 1 year unless noted otherwise below. In cases where the initial approval is authorized in months, 1 month is equal to 30 days.

Drug Policy

*Documentation Requirements:

The Company reserves the right to request additional documentation as part of its coverage determination process. The Company may deny reimbursement when it has determined that the drug provided or services performed were not medically necessary, investigational or experimental, not within the scope of benefits afforded to the member and/or a pattern of billing or other practice has been found to be either inappropriate or excessive. Additional documentation supporting medical necessity for the services provided must be made available upon request to the Company. Documentation requested may include patient records, test results and/or credentials of the provider ordering or performing a service. The Company also reserves the right to modify, revise, change, apply and interpret this policy at its sole discretion, and the exercise of this discretion shall be final and binding.

REFERENCES

1. Katz PO, Gerson LB, Vela MF. Diagnosis and management of gastroesophageal reflux disease. *Am J Gastroenterol*. 2013; 108:308–328.
2. AGA Institute. American Gastroenterological Association medical position statement on the management of gastroesophageal reflux disease. *Gastroenterology*. 2008;135:1383-1391.
3. Prilosec OTC® monograph. Available at: <http://www.prilosecotc.com/en-us/hcp/prilosec-otc-dosage>. Accessed on 17 January 2019.
4. Prevacid® 24HR Drug Facts. Available at: <http://www.prevacid24hr.com/index.jsp>. Accessed on 17 January 2019.
5. Zegerid OTC® [prescribing information]. Available at: <http://www.zegeridotc.com/zegeridotc/home/index.jspa>. Accessed on 17 January 2019.
6. Nexium® 24HR [prescribing information]. Available at: <https://www1.pfizerpro.com/hcp/nexium24hr/dosage-side-effects>. Accessed on 17 January 2019.
7. Prilosec® [prescribing information]. Willmington, DE: AstraZeneca, Inc; October 2016.
8. Prevacid® [prescribing information]. Deerfield, IL: Takeda Pharmaceuticals America, Inc; October 2016.
9. Protonix® [prescribing information]. Philadelphia, PA: Wyeth Laboratories; October 2016.
10. Aciphex® and Aciphex® Sprinkle™ [prescribing information]. Titusville, NJ; Janssen Pharmaceutica Inc., October 2016.
11. Nexium® [prescribing information]. Willmington, DE: AstraZeneca LP; October 2016.
12. Zegerid® [prescribing information]. San Diego, CA: Santarus, Inc.; October 2016.
13. Dexilant™ [prescribing information]. Deerfield, IL: Takeda Pharmaceuticals America, Inc.; October 2016.
14. Esomeprazole strontium delayed-release capsules [prescribing information]. Glasgow, KY: Amneal Pharmaceuticals; December 2014.
15. Larson C. Bioavailability and efficacy of omeprazole given orally and by nasogastric tube. *Dig Dis Sci*. 1996;41(3):475-479.
16. Sharma VK, Heinzlmann EJ, Steinberg EN, et al. Nonencapsulated, intact omeprazole granules effectively suppress intragastric acidity when administered via a gastrostomy. *Am J Gastroenterol*. 1997;92:848-851.
17. Chun AH. Lansoprazole: administration of the contents of a capsule dosage formulation through a nasogastric tube. *Clin Ther*. 1996;18(5):833-842.
18. Ying VWC, Kim SHH, Khan KJ, et al. Prophylactic PPI help to reduce marginal ulcers after gastric bypass surgery: a systemic review and meta-analysis of cohort studies. *Surg Endosc*. 2015;29:1018-1023.
19. Schulman AR, Chan W, Devery A, et al. Opened proton pump inhibitor capsules reduce time to healing compared with intact capsules for marginal ulceration following roux-en-Y gastric bypass. *Clin Gastroenterol Hepatol*. 2016 Oct. 20 [Epub ahead of print].
20. Drepper MD1, Spahr L, Frossard JL. Clopidogrel and proton pump inhibitors--where do we stand in 2012? *World J Gastroenterol*. 2012 May 14;18(18):2161-71.