

Drug Policy

Policy:	Sotyktu (deucravacitinib) oral tablets	Annual Review Date: 11/16/2023 Last Revised Date: 11/16/2023
----------------	-----------------------------------------------	-----------------------------------------------------------------------------------

OVERVIEW

Sotyktu, a tyrosine kinase 2 (TYK2) inhibitor, is indicated for treatment of moderate to severe **plaque psoriasis** in adults who are candidates for systemic therapy or phototherapy. Limitation of use: Sotyktu is not recommended in combination with potent immunosuppressants.

POLICY STATEMENT

This policy involves the use of **Sotyktu**. Prior authorization is recommended for pharmacy benefit coverage of **Sotyktu**. Approval is recommended for those who meet the conditions of coverage in the **Criteria and Initial/Extended Approval** for the diagnosis provided. **Conditions Not Recommended for Approval** are listed following the recommended authorization criteria. Requests for uses not listed in this policy will be reviewed for evidence of efficacy and for medical necessity on a case-by-case basis.

Because of the specialized skills required for evaluation and diagnosis of patients treated with **Sotyktu** as well as the monitoring required for adverse events and long-term efficacy, initial approval requires **Sotyktu** be prescribed by or in consultation with a physician who specializes in the condition being treated. All approvals for initial therapy are provided for the initial approval duration noted below; if reauthorization is allowed, a response to therapy is required for continuation of therapy unless otherwise noted below. **Sotyktu** is subject to the Inflammatory Conditions Care Value Program under pharmacy benefits.

RECOMMENDED AUTHORIZATION CRITERIA

Coverage of **Sotyktu** is recommended in those who meet the following criteria:

1. **Plaque Psoriasis.** Approve for the duration noted below if the patient meets ONE of the following (A or B):
 - A) **Initial Therapy.** Approve for if the patient meets the following criteria (i, ii, and iii):
 - i. Patient is \geq 18 years of age; AND
 - ii. Patient meets ONE of the following (a or b):
 - a) Patient has tried at least one traditional systemic agent for psoriasis for at least 3 months, unless intolerant;
OR
Note: Examples of one traditional systemic agent include methotrexate, cyclosporine, acitretin tablets, or psoralen plus ultraviolet A light (PUVA). An exception to the requirement for a trial of one traditional systemic agent for psoriasis can be made if the patient has already had a 3-month trial or previous

Drug Policy

intolerance to at least one biologic other than the requested drug. A biosimilar of the requested biologic does not count. Refer to [Appendix](#) for examples of biologics used for plaque psoriasis. A patient who has already tried a biologic for psoriasis is not required to “step back” and try a traditional systemic agent for psoriasis.

b) Patient has a contraindication to methotrexate, as determined by the prescriber; AND

iii. The medication is prescribed by or in consultation with a dermatologist.

B) Patient is Currently Receiving Sotyktu. Approve for duration noted below if patient meets the following criteria (i, ii, and iii):

i. Patient has been established on therapy for at least 90 days; AND

Note: A patient who has received < 90 days of therapy or who is restarting therapy is reviewed under criterion A (Initial Therapy).

ii. Patient experienced a beneficial clinical response, defined as improvement from baseline (prior to initiating the requested drug) in at least one of the following: estimated body surface area, erythema, induration/thickness, and/or scale of areas affected by psoriasis; AND

iii. Compared with baseline (prior to initiating the requested drug), patient experienced an improvement in at least one symptom, such as decreased pain, itching, and/or burning.

Initial Approval/ Extended Approval.

A) *Initial Approval:* 3 months (90 days)

B) *Extended Approval:* 1 year (365 days)

CONDITIONS NOT RECOMMENDED FOR APPROVAL

Sotyktu has not been shown to be effective, or there are limited or preliminary data or potential safety concerns that are not supportive of general approval for the following conditions. (Note: This is not an exhaustive list of Conditions Not Recommended for Approval).

- 1. Concurrent Use with other Biologics or with Targeted Synthetic Disease-Modifying Antirheumatic Drugs (DMARDs).** Data are lacking evaluating concomitant use of Sotyktu with another biologic or with a targeted synthetic DMARD for an inflammatory condition (see Appendix for examples). Combination therapy with biologics and/or biologics + targeted synthetic DMARDs has a potential for a higher rate of adverse effects and lack controlled trial data in support of additive efficacy. Note: This does NOT exclude the use of methotrexate (a traditional systemic agent used to treat psoriasis) in combination with Sotyktu.
- 2. Concurrent use with Other Potent Immunosuppressants, Including Methotrexate.** Co-administration with other potent immunosuppressive drugs has the risk of added immunosuppression and has not been evaluated.
- 3.** Coverage is not recommended for circumstances not listed in the Recommended Authorization Criteria. Criteria will be updated as new published data are available.

Drug Policy

Documentation Requirements:

The Company reserves the right to request additional documentation as part of its coverage determination process. The Company may deny reimbursement when it has determined that the drug provided or services performed were not medically necessary, investigational or experimental, not within the scope of benefits afforded to the member and/or a pattern of billing or other practice has been found to be either inappropriate or excessive. Additional documentation supporting medical necessity for the services provided must be made available upon request to the Company. Documentation requested may include patient records, test results and/or credentials of the provider ordering or performing a service. The Company also reserves the right to modify, revise, change, apply and interpret this policy at its sole discretion, and the exercise of this discretion shall be final and binding.

REFERENCES

1. Sotyktu™ tablets [prescribing information]. Princeton, NJ: Bristol Myers Squibb; September 2022.
2. Menter A, Strober BE, Kaplan DH, et al. Joint AAD-NPF guidelines of care for the management and treatment of psoriasis with biologics. *J Am Acad Dermatol.* 2019;80(4):1029-1072.
3. Nast A, Gisondi P, Ormerod AD, et al. European S3-Guidelines on the systemic treatment of psoriasis vulgaris – Update 2015 – Short version – EDF in cooperation with EADV and IPC. *J Eur Acad Dermatol Venereol.* 2015;29(12):2277-2294.

APPENDIX:

	Mechanism of Action	Examples of Inflammatory Indications*
Biologics		
Adalimumab SC Products (Humira®, biosimilars)	Inhibition of TNF	AS, CD, JIA, PsO, PsA, RA, UC
Cimzia® (certolizumab pegol SC injection)	Inhibition of TNF	AS, CD, nr-axSpA, PsO, PsA, RA
Etanercept SC Products (Enbrel®, biosimilars)	Inhibition of TNF	AS, JIA, PsO, PsA
Infliximab IV Products (Remicade®, biosimilars)	Inhibition of TNF	AS, CD, PsO, PsA, RA, UC
Simponi®, Simponi® Aria™ (golimumab SC injection, golimumab IV infusion)	Inhibition of TNF	SC formulation: AS, PsA, RA, UC
		IV formulation: AS, PJIA, PsA, RA
Actemra® (tocilizumab IV infusion, tocilizumab SC injection)	Inhibition of IL-6	SC formulation: PJIA, RA, SJIA
		IV formulation: PJIA, RA, SJIA
Kevzara® (sarilumab SC injection)	Inhibition of IL-6	RA
Orencia® (abatacept IV infusion, abatacept SC injection)	T-cell costimulation modulator	SC formulation: JIA, PSA, RA
		IV formulation: JIA, PsA, RA
Rituximab IV Products (Rituxan®, biosimilars)	CD20-directed cytolytic antibody	RA
Kineret® (anakinra SC injection)	Inhibition of IL-1	JIA [^] , RA
Stelara® (ustekinumab SC injection, ustekinumab IV infusion)	Inhibition of IL-12/23	SC formulation: CD, PsO, PsA, UC
		IV formulation: CD, UC
Siliq™ (brodalumab SC injection)	Inhibition of IL-17	PsO
Cosentyx® (secukinumab SC injection)	Inhibition of IL-17A	AS, ERA, nr-axSpA, PsO, PsA
Taltz® (ixekizumab SC injection)	Inhibition of IL-17A	AS, nr-axSpA, PsO, PsA
Ilumya™ (tildrakizumab-asmn SC injection)	Inhibition of IL-23	PsO
		SC formulation: CD, PSA, PsO

This document is subject to the disclaimer found at <https://www.medmutual.com/For-Providers/Policies-and-Standards/CorporateMedicalDisclaimer.aspx> and is subject to change. <https://www.medmutual.com/For-Providers/Policies-and-Standards/Prescription-Drug-Resources.aspx>

Drug Policy

Skyrizi® (risankizumab-rzaa SC injection, risankizumab-rzaa IV infusion)		IV formulation: CD
Tremfya™ (guselkumab SC injection)	Inhibition of IL-23	PsO
Entyvio™ (vedolizumab IV infusion)	Integrin receptor antagonist	CD, UC
Oral Therapies/Targeted Synthetic DMARDs		
Otezla® (apremilast tablets)	Inhibition of PDE4	PsO, PsA
Cibinco™ (abrocitinib tablets)	Inhibition of JAK pathways	AD
Olumiant® (baricitinib tablets)	Inhibition of JAK pathways	RA
Rinvoq® (upadacitinib extended-release tablets)	Inhibition of JAK pathways	AD, AS, nr-axSpA, RA, PsA, UC
Sotyktu™ (deucravacitinib tablets)	Inhibition of TYK2	PsO
Xeljanz® (tofacitinib tablets)	Inhibition of JAK pathways	RA, PJIA, PsA, UC
Xeljanz® XR (tofacitinib extended-release tablets)	Inhibition of JAK pathways	RA, PsA, UC

* Not an all-inclusive list of indications (e.g., oncology indications and rare inflammatory conditions are not listed). Refer to the prescribing information for the respective agent for FDA-approved indications; SC – Subcutaneous; TNF – Tumor necrosis factor; AS – Ankylosing spondylitis; CD – Crohn’s disease; JIA – Juvenile idiopathic arthritis; PsO – Plaque psoriasis; PsA – Psoriatic arthritis; RA – Rheumatoid arthritis; UC – Ulcerative colitis; nr-axSpA – Non-radiographic axial spondyloarthritis; IV – Intravenous, PJIA – Polyarticular juvenile idiopathic arthritis; IL – Interleukin; SJIA – Systemic juvenile idiopathic arthritis; ^ Off-label use of Kineret in JIA supported in guidelines; ERA – Enthesitis-related arthritis; DMARD – Disease-modifying antirheumatic drug; PDE4 – Phosphodiesterase 4; JAK – Janus kinase; AD – Atopic dermatitis; TYK2 – Tyrosine kinase 2.